

Fig. 1

Fig. 1A

Fig. 1B

Fig. 1C




Fig. 1D

Fig. 1E

Fig. 2



Fig. 2A




Fig. 2B

Fig. 3

Recovery from 48 hour exposure to
tunicamycin (BAMEC)




Fig. 3A

Recovery from 72 hour exposure to
tunicamycin (BAMEC)




Fig. 3B

Fig. 4




Fig. 4A




Fig. 4B

Fig. 5




Fig. 6

Fig. 6A




Fig. 6B




Fig. 6

Fig. 6A




Fig. 6B




Fig. 7

Fig. 7A




Fig. 7B




Fig. 7C




Fig. 7

Fig. 7D




Fig. 7E




Fig. 7F




Fig. 8




Fig. 9

Fig. 9A



Fig. 9B



Fig. 9C




Fig. 10





Fig. 10A

Fig. 10B

Flow cytometry histogram showing cell population distribution. The x-axis is labeled 'FL2-H' and the y-axis is labeled 'SSC-H'. The plot shows a single peak of cells, with a gate labeled 'M1' indicating the selected population.

Fig. 10





Fig. 10C




Fig. 10D

Fig. 11





Fig. 11B

Fig. 11A

Fig. 11C

Fig. 11




Fig. 11C

Fig. 11D

Fig. 12




Fig. 13

Fig. 13A




Fig. 13E




Fig. 13F




Fig. 13

Fig. 13B




Fig. 13C




Fig. 13D




Fig. 13

Fig. 13





Fig. 13H

Fig. 13G

Fig. 14

